

## Blood-clotting protein linked to cancer and septicaemia

Research could lead to new therapies

**Heidelberg, 4 February 2011** - In our not-so-distant evolutionary past, stress often meant imminent danger, and the risk of blood loss, so part of our body's stress response is to stock-pile blood-clotting factors. Scientists in the Molecular Medicine Partnership Unit (MMPU), a collaboration between the European Molecular Biology Laboratory (EMBL) in Heidelberg, Germany, and the University of Heidelberg Medical Centre, have discovered how stressed cells boost the production of the key blood-clotting factor, thrombin. Their work, published today in *Molecular Cell*, shows how cancer cells may be taking advantage of this process, and opens new possibilities for fighting back, not only against cancer but also against septicaemia, where increased blood clotting is still one of the leading causes of death.

Blood clots tend to form more often in the veins of people with cancer, a syndrome first described almost 150 years ago by French physician Armand Trousseau. In recent years, doctors have also come to realise that people with activated blood coagulation are more likely to develop cancer. Accordingly, recent studies have shown that anti-coagulants may help treat and prevent cancer, but exactly how blood-clotting and cancer progression are linked was unclear – until now.

"For the 1st time, we have something in hand that might explain this enigmatic relationship between enhanced pro-coagulatory activities and the outcome of cancer," says Sven Danckwardt, who carried out the research within the MMPU.

The amount of thrombin our cells produce is controlled by two sets of proteins: proteins that slow thrombin production, and proteins that speed it up. Both types of protein act by binding to the cellular machinery that synthesises thrombin, and, under normal conditions, the production-slowing proteins keep thrombin levels low. But Danckwardt and colleagues discovered that, when our cells come under stress from inflammation, another protein, called p38 MAPK, reacts by adding a chemical tag to those production-slowing proteins. That tag makes it harder for the production-slowing proteins to bind to the thrombin-synthesising machinery, allowing the

proteins that speed up production to take over. So inflammation caused by cancer could lead to increased thrombin levels and, as thrombin is a blood-clotting agent, this could explain why cancer patients are more likely to suffer from blood-clots. The scientists believe this new mechanism of gene regulation may apply to other genes, too.

"Knowing the exact molecules involved, and how they act, has implications for treatment, especially as drugs that inhibit p38 MAPK are already being tested in clinical studies for other conditions," says Matthias Hentze, Associate Director of EMBL and co-director of MMPU, adding: "those drugs could be good candidates for potential cancer or septicaemia therapies."

The Heidelberg scientists found that p38 MAPK also influences thrombin production during septicaemia. Also known as blood poisoning, septicaemia occurs when bacteria or other pathogens enter the bloodstream, leading to widespread infection and to blood-clotting problems. When they analysed liver samples taken from septicaemic mice and from cancer patients, the scientists discovered that thrombin production increases in response both to widespread inflammation during septicaemia and to localized inflammation at the tumour's invasion front, where cancer cells are spreading into nearby tissue.

Aside from its role as a blood-clotting agent, thrombin is also involved in creating new blood vessels, and it is able to degrade the extracellular matrix that keeps cells together. So it's possible that the cancer cells are increasing thrombin production to help the tumour spread, by making it easier to invade healthy tissue and creating blood vessels to supply the new tumour cells. This, the researchers believe, could explain why people with blood-clotting problems seem to have a higher risk of developing cancer.

"This study shows the benefits of partnerships like the MMPU, which bridge the gap between clinical and basic research," Andreas Kulozik from the University of Heidelberg Medical Centre, co-director of MMPU, concludes. ●

### Source Article

Danckwardt, S., Gantert, A.-S., Macher-Goeppinger, S., Probst, H.C., Gentzel, M., Wilm, M., Gröne, H.-J., Schirmacher, P., Hentze, M.W., Kulozik, A.E. p38 MAPK controls prothrombin expression by regulated RNA 3'end processing. *Molecular Cell*, 4 February 2011

### Contact:

Sonia Furtado, EMBL Press Officer, Heidelberg, Germany, Tel: +49 6221 387 8263, [www.embl.org](http://www.embl.org), [sonia.furtado@embl.de](mailto:sonia.furtado@embl.de)

## About EMBL

The European Molecular Biology Laboratory is a basic research institute funded by public research monies from 20 member states (Austria, Belgium, Croatia, Denmark, Finland, France, Germany, Greece, Iceland, Ireland, Israel, Italy, Luxembourg, the Netherlands, Norway, Portugal, Spain, Sweden, Switzerland and the United Kingdom) and associate member state Australia. Research at EMBL is conducted by approximately 85 independent groups covering the spectrum of molecular biology. The Laboratory has five units: the main Laboratory in Heidelberg, and Outstations in Hinxton (the European Bioinformatics Institute), Grenoble, Hamburg, and Monterotondo near Rome. The cornerstones of EMBL's mission are: to perform basic research in molecular biology; to train scientists, students and visitors at all levels; to offer vital services to scientists in the member states; to develop new instruments and methods in the life sciences and to actively engage in technology transfer activities. Around 190 students are enrolled in EMBL's International PhD programme. Additionally, the Laboratory offers a platform for dialogue with the general public through various science communication activities such as lecture series, visitor programmes and the dissemination of scientific achievements.

### *Policy regarding use*

*EMBL press and picture releases including photographs, graphics, movies and videos are copyrighted by EMBL. They may be freely reprinted and distributed for non-commercial use via print, broadcast and electronic media, provided that proper attribution to authors, photographers and designers is made. High-resolution copies of the images can be downloaded from the EMBL web site: [www.embl.org](http://www.embl.org)*